

GRAVITATIONAL WAVE RECOIL OSCILLATIONS OF BLACK HOLES: IMPLICATIONS FOR UNIFIED MODELS OF ACTIVE GALACTIC NUCLEI

S. KOMOSSA

Max-Planck-Institut für extraterrestrische Physik, Postfach 1312, 85741 Garching, Germany; skomossa@mpe.mpg.de

DAVID MERRITT

Center for Computational Relativity and Gravitation and Department of Physics, Rochester Institute of Technology, Rochester, NY 14623, USA; merritt@astro.rit.edu

Draft version November 7, 2008

ABSTRACT

We consider the consequences of gravitational wave recoil for unified models of active galactic nuclei (AGNs). Spatial oscillations of supermassive black holes (SMBHs) around the cores of galaxies following gravitational wave (GW) recoil imply that the SMBHs spend a significant fraction of time off-nucleus, at scales beyond that of the molecular obscuring torus. Assuming reasonable distributions of recoil velocities, we compute the off-core timescale of (intrinsically type-2) quasars. We find that roughly one-half of major mergers result in a SMBH being displaced beyond the torus for a time of $10^{7.5}$ yr or more, comparable to quasar activity timescales. Since *major* mergers are most strongly affected by GW recoil, our results imply a deficiency of type 2 quasars in comparison to Seyfert 2 galaxies. Other consequences of the recoil oscillations for the observable properties of AGNs are also discussed.

Subject headings: galaxies: active – galaxies: evolution – quasars: general

1. INTRODUCTION

Gravitational waves emitted anisotropically during gravitational collapse carry away linear momentum. As a result the center of mass of the collapsing object recoils (Peres 1962; Bekenstein 1973). Configurations of coalescing spinning black holes can result in recoil velocities of hundreds to thousands of km s^{-1} (e.g., Campanelli et al. 2007; González et al. 2007; Herrmann et al. 2007; Pollney et al. 2007; Tichy & Marronetti 2007; Brüggemann et al. 2008; Dain et al. 2008), scaling to a maximum of $\sim 4000 \text{ km s}^{-1}$ (Campanelli et al. 2007; Baker et al. 2008) for maximally spinning equal-mass binaries with anti-aligned spins in the orbital plane, and as large as $\sim 10^4 \text{ km s}^{-1}$ in hyperbolic encounters (Healy et al. 2008). Kicks large enough to remove SMBHs from galaxies have potentially far-reaching consequences for SMBH and galaxy assembly, and predict interstellar and intergalactic quasars (e.g., Madau et al. 2004; Merritt et al. 2004; Madau & Quataert 2004; Haiman 2004; Boylan-Kolchin et al. 2004; Libeskind et al. 2006; Loeb 2007; Schnittman 2007; Volonteri 2007; Gualandris & Merritt 2008; Volonteri & Madau 2008, Kornreich & Lovelace 2008).

Komossa et al. (2008) reported the detection of a recoil candidate with a projected kick velocity of 2650 km s^{-1} . The quasar SDSSJ092712.65+294344.0 shows three unusual emission-line systems, including a kinematically offset broad-line region (BLR) and a system of atypically narrow emission lines which lack the usual ionization stratification – two key signatures of kicks. Apart from spectroscopic signatures (Merritt et al. 2006; Bonning et al. 2007), offset quasars could be detected for instance by their temporarily flaring accretion disks (Shields & Bonning 2008; Lippai et al. 2008; Schnittman & Krolik 2008), by tidal disruption flares from the bound (and unbound) population of stars and episodic fuelling from

stellar mass loss (Komossa & Merritt 2008), and via the bound compact star cluster itself (Merritt et al. 2008, O’Leary & Loeb 2008). One key consequence of gravitational wave recoil is long-lasting oscillations of the SMBH about the galaxy core, implying that SMBHs may spend as long as 10^{6-9} yrs off-nucleus with an amplitude of parsecs or kiloparsecs, depending on kick velocity and galaxy structural parameters (Merritt et al. 2004; Madau & Quataert 2004; Gualandris & Merritt 2008; MQ04 and GM08 hereafter). In this Letter, we consider consequences of these “recoil oscillations” for unified models of active galactic nuclei (AGNs).

According to unified models, AGNs are intrinsically similar, but their appearance depends strongly on the line of sight of the observer toward the “central engine” (reviews by Antonucci 1993; Elitzur 2007). Along certain sightlines, a dusty torus consisting of molecular clouds blocks the observer’s view, hiding some core components, especially the BLR, which therefore is only directly visible in “type 1” AGNs. The unified model has been very successful in explaining observed properties of AGNs including the presence of hidden BLRs detected in polarized light (e.g., Antonucci & Miller 1985; Zakamska et al. 2005), and has been corroborated by recent imaging and spectroscopy of the torus (e.g., Jaffe et al. 2004; Siebenmorgen et al. 2005).

If SMBHs and the BLRs bound to them spend a significant time displaced from the nucleus, this will have profound consequences for obscuration-based unified models of AGNs. This statement holds true whether the obscuration originates in parsec-scale molecular tori (Antonucci 1993), or compact star-forming regions (Levenson et al. 2001), or is due to absorption associated with the host galaxy itself, which may work on typical scales of $\sim 100 \text{ pc}$ (Maiolino & Rieke 1995). Here we argue that a significant fraction of the quasar population is ex-

pected to be in a regime such that the SMBH and the BLR bound to it is displaced beyond the obscuring region, implying a deficiency of type 2 (obscured) AGNs among quasars.

2. RELEVANT SPATIAL SCALES IN THE AGN CORE

We first compare three characteristic spatial scales: the kick radius, the BLR size, and the torus size. After the kick, matter remains bound to the SMBH within a region whose radius r_k is given by

$$r_k = \frac{GM_{\text{BH}}}{v_k^2} \approx 0.43 \left(\frac{M_{\text{BH}}}{10^8 M_\odot} \right) \left(\frac{v_k}{10^3 \text{ km s}^{-1}} \right)^{-2} \text{ pc} \quad (1)$$

where v_k is the kick velocity (e.g., Merritt et al. 2006). The size r_{BLR} of the BLR of AGNs has been determined from reverberation mapping (Peterson 2007), and scales with AGN luminosity as

$$r_{\text{BLR}} = 0.1 \left(\frac{\lambda L_\lambda(5100\text{\AA})}{10^{45} \text{ erg s}^{-1}} \right)^{0.69} \text{ pc} \quad (2)$$

(Kaspi et al. 2005) where the luminosity at 5100\AA , $\lambda L_\lambda(5100\text{\AA}) \simeq 0.1 L$ and L is the AGN bolometric luminosity. The size of the molecular torus is still poorly constrained from observations. Recent measurements of dusty gas in the Seyfert galaxies NGC 1068 and Circinus suggest an extent of 2-3 pc (Jaffe et al. 2004; Davies et al. 2007; Tristram et al. 2007). It is reasonable to assume that the *inner* edge of the torus, $r_{\text{tor,in}}$, is beyond the dust sublimation radius (e.g., Netzer 1990; Nenkova et al. 2008) which is given by

$$r_{\text{tor,in}} \gtrsim 0.4 \left(\frac{L}{10^{45} \text{ erg s}^{-1}} \right)^{1/2} \left(\frac{1500\text{K}}{T_{\text{sub}}} \right)^{2.6} \text{ pc} \quad (3)$$

with an outer radius likely not much larger than $\sim 20 r_{\text{tor,in}}$ (Elitzur 2007), and where T_{sub} is the dust sublimation temperature. Comparison of these three relations shows that a large fraction of the BLR remains bound to the recoiling hole, while structures of the size of the torus or larger will typically be left behind. Oscillation amplitudes of $\gtrsim 10$ -20 pc will therefore move the SMBH beyond the torus scale, except for the highest SMBH masses, where part of the torus will remain bound.

3. OSCILLATION AMPLITUDES AND OSCILLATION DURATIONS IN QUASARS

We base our discussion on the N -body simulations of GM08, who computed SMBH trajectories after GW recoil. Key model parameters are the SMBH mass M_{BH} and galaxy mass $M_{\text{gal}} = 10^3 M_{\text{BH}}$, the kick velocity v_k , and the galaxy structural parameters. We concentrate here on the mass range of SMBHs that is typical for the bulk of quasars ($M_{\text{BH}} \approx 10^8 - \text{few } 10^9 M_\odot$). Massive inactive elliptical galaxies and host galaxies of luminous quasars typically have effective radii R_e on the order of a few to ~ 10 kpc. Here, we adopt the relation between R_e and M_{gal} defined by the nine luminous ($-22 \lesssim M_V \lesssim -24$) quasar host galaxies in the sample of Wolf & Sheinis (2008), i.e. $\log(R_e/\text{kpc}) \approx -5.61 + 0.55 \log(M_{\text{gal}}/M_\odot)$. The models of GM08 on which we base our discussion had post-kick core radii r_c consistent with the range defined by the brightest E-galaxies with resolved cores (Ferrarese et al. 2006), i.e. $0.01 \lesssim r_c/R_e \lesssim 0.03$. We focus here on

the spherical ‘‘A1’’ galaxy models from GM08; scaling of the N -body results to physical units was done following their equation (4). Dark matter halos were ignored.

Figure 1a shows trajectories of kicked SMBHs scaled to a galaxy with SMBH mass $5 \times 10^8 M_\odot$. Kick velocities were $v_k/v_{\text{esc}} = 0.3, 0.5, 0.7, 0.9$, corresponding to $v_k = (360, 590, 830, 1070) \text{ km s}^{-1}$ for a central escape velocity v_{esc} of 1185 km s^{-1} of our galaxy model. As discussed by GM08, SMBH oscillations continue well beyond the time (‘‘Phase I’’) that would be predicted by applying Chandrasekhar’s dynamical friction formula assuming a fixed galaxy core (e.g. MQ04; Blecha & Loeb 2008). The time of onset of these long-term, or ‘‘Phase II’’, oscillations is indicated by the open circles in Figure 1a. Including the effect of the Phase II oscillations, Figure 1a shows that the SMBH’s motion persists for more than $\sim 10^7$ yr if $v_k \gtrsim 450 \text{ km s}^{-1}$.

4. COMPARISON WITH OBSERVATIONS

4.1. Unified models and type1/type2 fractions

About 70% of the nearby Seyfert galaxies are type 2 (e.g., Schmitt et al. 2001), i.e. they lack a BLR in their (unpolarized) optical spectra. The fraction of the high-luminosity equivalents, type 2 quasars, is less well known and still subject to a number of selection effects (e.g. Halpern & Moran 1998; Reyes et al. 2008), even though studies generally indicate a deficiency of type 2 quasars. While in past studies type 2 quasars were observationally very rare or absent, significant numbers have been found in recent X-ray and optical surveys (e.g., Norman et al. 2002; Zakamska et al. 2005; Brusa et al. 2007). There is a systematic trend such that the fraction of type 2 sources, or equivalently the amount of X-ray absorption, decreases with increasing source luminosity (e.g., Simpson 2005; Barger et al. 2005; Reyes et al. 2008; Hasinger 2008), while the simplest possible version of the unified model would imply a constant type1/type2 ratio. Models have been proposed which account for this luminosity dependence, e.g. by invoking changes in the properties of the obscurer as L increases (e.g., Nenkova et al. 2008; Ballantyne 2008). Recoil oscillations inevitably affect the numbers of obscured versus unobscured sources and therefore have potentially profound implications for unified models. How do they affect the ratio of type1/type2 quasars in comparison to the number of type1/type2 Seyfert galaxies? There is increasing evidence that quasar activity is powered by major mergers while Seyfert activity may have other triggers including bars, minor mergers, or random accretion of molecular clouds (e.g., Sanders et al. 1988; Urrutia et al. 2008; Hopkins et al. 2008; Hasinger 2008, and references therein). Recoil oscillations have highest amplitudes in major mergers, and we may therefore expect that the frequency and properties of (type 2) quasars are strongly affected by recoil oscillations. Can this explain the relative scarcity of type 2 quasars in comparison with type 2 Seyferts, and the more general trend that type2-ness (X-ray absorption) decreases with luminosity?

4.2. Rate estimates

Since a large fraction of quasars are believed to be triggered by major mergers, in order to estimate the fraction of quasars that occur at or above a given kick velocity,

we carried out rate estimates relevant for major mergers with random spin distributions (Campanelli et al. 2007; Schnittman & Buanano 2007; Baker et al. 2008; note that the actual kick velocities could be smaller in gas-rich systems if the mechanism discussed by Bogdanović et al. 2007 is at work, but see Sect. 3 of Schnittman & Krolik 2008). We first updated these previous estimates, based on the recent kick formula of Baker et al. (2008; essentially identical results were obtained using the Lousto & Zlochower 2008 version of the kick formula), and assuming random orientations of the spin vectors of both SMBHs and a distribution of SMBH mass ratios in the range $0.3 \leq q \leq 1.0$ where $q \equiv m_2/m_1, m_2 \leq m_1$, relevant for major mergers. SMBH spins were drawn from a distribution such that $a_1 \leq a \leq a_2$ with $a \equiv S/m^2$ the dimensionless spin, with $a_1 = 0.5, a_2 = 0.9$. The distributions of m and a were assumed to be uniform in the logarithm between these limits. For this base model, we find that about 50% of major mergers have kick velocities above 500 km/s. At or above these kick velocities, *total* oscillation timescales start to be on the order of quasar lifetimes, which are about 10^{7-8} yr (e.g., Kauffmann & Haehnelt 2000; Yu & Tremaine 2002; Hopkins et al. 2005; see review by Martini 2004).

How much of the total quasar population is ultimately affected by recoil oscillations then depends on (1) the time t_{vis} the kicked SMBH+BLR spends *beyond* the obscuring torus (i.e., at distances greater than $r_{\text{torus}} \approx r_{\text{tor,out}}$ from the galaxy center (eqn. 3)) and therefore appears as “type 1” rather than “type 2”, in comparison to (2) the total quasar lifetime. We computed t_{vis} on a grid in r_{torus} for each of the nine N -body trajectories in GM08 ($0.1 \leq v_k/v_{\text{esc}} \leq 0.9$). Given arbitrary values of v_k, M_{gal} and r_{torus} , the value of t_{vis} was then computed via interpolation between the nine discrete kick velocities of GM08.

Figure 1b shows, for our base model, the fraction f of kicked SMBHs that spend more than 3×10^7 yr at $r > r_{\text{min}}$, for a range of $(M_{\text{gal}}, r_{\text{min}})$ values. For $r_{\text{min}} \approx r_{\text{tor,out}}$, this fraction is $\sim 0.5 \pm 0.1$ with a weak dependence on galaxy mass. We conclude that *a significant fraction of mergers would result in recoiling SMBHs that spend of order a quasar lifetime above the obscuring torus* and would therefore appear as type 1 quasars even if “intrinsically” type 2. We examined the robustness of these results under different assumptions about the pre-recoil distributions of spins and masses. Maximally spinning, equal-mass SMBH mergers affect a fraction of up to $f \sim 0.75$ of the population (same assumptions as above), while major mergers with intermediate SMBH spins of $a = 0.3$ imply $f \sim 0.05 - 0.15$.¹

If we assume that the type1/type2 fraction of quasars is intrinsically (i.e., in the absence of recoils) the same as in Seyferts ($\sim 70\%$ type 2, $\sim 30\%$ type 1), we predict that a fraction $\sim 50\%$ of these type2s will appear instead as type 1s at any given time. Several factors affect these estimates: idealizations in the galaxy models used in the N -body simulations, uncertainties in the distribution of masses and spins as discussed above, uncer-

tainties especially in the thickness of the torus in the most massive galaxies, and various selection effects in the measurements of type1/type2 ratios in dependence of SMBH mass. We also recall that if equations (1) and (3) strictly hold, in the most massive quasars a fraction of the torus will remain bound to the recoiling SMBH so absorption/extinction would not fall to zero. Finally, we note that if the quasars’ *total* lifetime is actually composed of several shorter merger episodes on the order of 10^6 yr each, even recoil oscillations with velocities as small as ~ 200 km s⁻¹ would affect a large fraction of the quasar population.

4.3. Implications

So far, we have distinguished between Seyfert galaxies (low-mass SMBHs) and quasars (high-mass SMBHs). Can we also reproduce the observed trend (Sect. 4.1) that obscuration fraction decreases with quasar luminosity (i.e., mass)? In our base model, the dependence of mean oscillation timescale on galaxy mass is weak. However, since the likelihood of a major merger (as opposed to other types of fuelling) is believed to increase strongly with AGN luminosity (galaxy mass), the observed trend should arise naturally, since the most luminous AGNs are increasingly likely to be triggered by major mergers.

If a fraction of all quasars is recoiling at any given time, we should see the corresponding BLR emission-line velocity shifts v_{obs} in a fraction of the type 1 quasars. Bonning et al. (2007; B07 hereafter) set limits on the fraction of emission line velocity shifts observed in a sample of SDSS quasars. In our model, the majority ($\gtrsim 75\%$) of kicked SMBHs remain bound to the galaxy and so their velocities quickly drop *below the initial value of v_k* ; most of the time they would therefore be observed with a velocity that is much smaller than v_k .

We carried out Monte-Carlo simulations to check the consistency of our recoil model with the B07 limits. Our simulations were designed to crudely mimic the properties of a sample of quasars selected to exhibit both broad and narrow emission lines, as in the B07 sample. We adopted a uniform logarithmic distribution of galaxy masses, $11 \leq \log_{10}(M_{\text{gal}}/M_{\odot}) \leq 12.5$, and a fixed quasar lifetime of $t_{\text{qso}} = 3 \times 10^7$ yr; kicks were assumed to have occurred at times that were distributed uniformly and randomly between the epoch of observation and a time t_{qso} earlier. The position and velocity of the SMBH at the time of observation was extracted from the appropriate N -body model after scaling to physical units using the galaxy mass. If the distance of the SMBH from the galaxy center exceeded the outer torus radius, its radial velocity v_{obs} was added to the cumulative distribution (assuming a random direction for the recoil); this velocity was identified with the measured velocity offset of the BLR gas in the B07 galaxies. A fraction 30% of SMBHs (the “true” type 1 population) with $r < r_{\text{tor,out}}$ were assumed to have visible BLRs and so were included in the accounting. Figure 1c shows that the predicted fraction of objects with large ($\gtrsim 100$ km s⁻¹) velocity shifts is \sim an order of magnitude smaller than would be inferred from the unmodified distribution of kick velocities; this is due to the deceleration that occurs as the SMBH moves through the galaxy.

B07 found a maximal fraction of $f_{500}=0.04$ quasars with velocity shifts above $v_{\text{obs}} = 500$ km s⁻¹, and a frac-

¹ We note that galaxy triaxiality, not included in the current models, might extend the oscillation timescales by a factor of several (e.g. Vicari et al. 2007; see also MQ04 and Merritt et al. 2004); inhomogeneities of the host galaxy after merging would likely have the same effect.

tion $f_{1000}=0.0035$ with shifts above $v_{\text{obs}} = 1000 \text{ km s}^{-1}$. These limits are consistent with our baseline model for the kick distribution, particularly if we impose an upper limit to the galaxy/SMBH separation at which a recoiling SMBH would be spectroscopically identified with its host galaxy (shown as the dotted lines in Figure 1c, assuming $r_{\text{max}} = 10 \text{ kpc}$).

Finally, we note that recoil oscillations will also have a number of other observable consequences. They will affect the X-ray background and its modeling since a fraction of sources will be unobscured at any given time. In particular, small amplitude oscillations of the order the torus size will affect the ratio of Compton-thin to Compton-thick sources, and could lead to measurable variability in the absorption and extinction of AGN spectra once the recoiling SMBH passes the individual clouds making up the torus. A number of interesting effects are related to the torus itself: (1) A recoiling SMBH with a bound gas disk that passes through the dense torus (rather than moving perpendicular to it) might cause local shocks, heating, and temporary X-ray emission. (2) During the long-lived ‘‘Phase II’’ oscillations, when the SMBH oscillation amplitude is on the torus scale, the SMBH might efficiently accrete from the dense molecular gas at *each* turning point, causing repeated flares of radiation. Such flares would locally destroy the dust, while photoionization of the dense surrounding gas would produce a strong emission-line response which would not

only help in identifying recoils but could also be used as a new probe of the properties of the torus itself.² (3) Torus radii are roughly equal to SMBH gravitational influence radii, so ejection of the SMBH might lead to temporary expansion of the torus since the mass holding it in place is suddenly removed. Isotropic expansion would not affect the column density along the line of sight, seen from the very center. These effects will be addressed in more detail in forthcoming work.

In summary, we have shown that timescales of recoil oscillations are in an interesting regime where they can potentially affect a significant fraction of the quasar population. Details of the predictions still depend on uncertainties in the observed numbers of AGN types, quasar lifetimes, torus properties, structural parameters of luminous quasar host galaxies, and SMBH spin distributions on the one hand, and on modelling recoil in realistic non-spherical or non-axially-symmetric galaxies on the other hand. Knowledge of recoil oscillation timescales and amplitudes is also critical for modelling AGN evolution, delays between starburst and AGN activity after merger, and the cosmic X-ray background.

DM was supported by grants AST-0807910 (NSF) and NNX07AH15G (NASA). We thank A. Gualandris for assistance in extracting data from the N -body simulations.

² While strong emission-line variability in response to an X-ray flare has recently been observed, this event is more likely inter-

preted in terms of stellar tidal disruption (Komossa et al. 2008b).

REFERENCES

- Antonucci, R.R.J. 1993, *ARA&A*, 31, 473
 Antonucci, R.R.J. & Miller, J.S., 1985, *ApJ*, 297, 621
 Baker, J.G., et al. 2008, *ApJ*, 682, L29
 Ballantyne, D.R. 2008, *ApJ*, in press, arXiv:0806.2863
 Barger, A.J., et al. 2005, *AJ*, 129, 578
 Bekenstein, J.D. 1973, *ApJ*, 183, 657
 Blecha, L., & Loeb, A. 2008, *MNRAS*, in press
 Bogdanović, T., et al. 2007, *ApJ*, 661, L147
 Bonning, E.W., et al. 2007, *ApJ*, 666, L13 (B07)
 Boylan-Kolchin, M., et al. 2004, *ApJ*, 613, L37
 Brüggmann, B., et al. 2008, *Phys. Rev. D.*, 77, 124047
 Brusa, M., et al. 2007, *ApJS*, 172, 353
 Campanelli, M., et al. 2007, *ApJ*, 659, L5
 Dain, S., et al. 2008, *Phys. Rev. D.*, 74, 024039
 Davies, R., et al. 2007, *ASPC*, 373, 639
 Elitzur, M. 2007, *ASPC*, 373, 415
 Ferrarese, L., et al. 2006, *ApJS*, 164, 334
 Gonzáles, J.A., et al. 2007, *Phys. Rev. Lett.*, 98, 231101
 Gualandris, A., & Merritt, D. 2008, *ApJ*, 678, 780 (GM08)
 Haiman, Z. 2004, *ApJ*, 613, 36
 Halpern, J., & Moran, E.C., 1998, 494, 194
 Hasinger, G., 2008, *A&A*, in press, arXiv:0808.0260
 Healy, J., et al. 2008, arXiv:0807.3292v1
 Herrmann, F., et al. 2007, *Phys. Rev. D*, 76, 084032
 Hopkins, P., et al. 2005, *ApJ*, 625, L71
 Hopkins, P., et al. 2008, *ApJS*, 175, 356
 Jaffe, W., et al. 2004, *Nature*, 429, 47
 Kaspi, S., et al. 2005, *ApJ*, 629, 61
 Kauffmann, G., & Haehnelt, M., 2000, *MNRAS*, 311, 576
 Komossa, S., Zhou H., & Lu H., 2008, *ApJ*, 678, L81
 Komossa, S., et al. 2008b, *ApJ*, 678, L13
 Komossa, S., & Merritt, D., 2008, *ApJ*, 683, L21
 Kornreich, D.A., & Lovelace, R.V.E., 2008, *ApJ*, 681, 104
 Levenson, N., et al. 2001, *ApJ*, 557, 54
 Libeskind, N.L., et al. 2006, *MNRAS*, 368, 1381
 Lippai, Z., et al. 2008, *ApJ*, 676, L5
 Loeb, A. 2007, *Phys. Rev. Lett.*, 99, 041103
 Lousto, C.O., & Zlochower, Y., 2008, arXiv:0805.0159
 Madau, P., et al. 2004, *ApJ*, 604, 484
 Madau, P., & Quataert, E. 2004, *ApJ*, 606, L17 (MQ04)
 Maiolino, R., & Rieke, G.H., 1995, *ApJ*, 454, 95
 Martini, P., 2004, in: *Carnegie Observatory Astrophysics Series 1*, L. Ho (ed.), 169
 Merritt, D., et al. 2004, *ApJ*, 607, L9
 Merritt, D., et al. 2006, *MNRAS*, 367, 1746
 Merritt, D., et al., 2008, *ApJ*, submitted, arXiv:0809.5046
 Nenkova, M., et al. 2008, *ApJ*, in press, arXiv:0806.0512
 Netzer, H. 1990, in: *Saas Fee Lecture Notes*, 20, 57
 Norman, C., et al. 2002, *ApJ*, 57, 218
 Peres, A. 1962, *Phys. Rev.*, 128, 2471
 O’Leary, R., & Loeb, A. 2008, submitted, arXiv:0809.4262
 Peterson, B.M. 2007, *ASPC*, 373, 3
 Pollney, D., et al. 2007, *Phys. Rev. D*, 76, 124002
 Reyes, R., et al. 2008, *AJ*, submitted, arXiv:0801.1115
 Sanders, D., et al. 1988, *ApJ*, 325, 74
 Schmitt, H.R., et al. 2001, *ApJ*, 555, 663
 Schnittman, J.D. 2007, *ApJ*, 667, L133
 Schnittman, J.D., & Buanano, A. 2007, *ApJ*, 662, L63
 Schnittman, J.D., & Krolik, J.H. 2008, *ApJ*, 684, 835
 Shields, G.A., & Bonning, E.W. 2008, *ApJ*, 682, 758
 Siebenmorgen, R., et al. 2005, *A&A*, 436, L5
 Simpson, C. 2005, *MNRAS*, 360, 565
 Tichy, W., & Marronetti, P. 2007, *Phys. Rev. D*, 76, 061502
 Tristram, K.R.W., et al. 2007, *A&A*, 474, 837
 Urrutia, T., et al. 2008, *ApJ*, 674, 80
 Vicari, A., et al. 2007, *ApJ*, 662, 797
 Volonteri, M. 2007, *ApJ*, 663, L5
 Volonteri, M., & Madau, P. 2008, *ApJ*, in press
 Wolf, M.J., & Sheinis, A.I. 2008, *AJ*, 136, 1587
 Yu, Q., & Tremaine, S., 2002, *MNRAS*, 335, 965
 Zakamska, N.L., et al. 2005, *AJ*, 129, 1212




FIG. 1.— (a) Trajectories of kicked SMBHs in N -body models of a galaxy (based on GM08) with a SMBH mass of $M_{\text{BH}} = 5 \times 10^8 M_{\odot}$, galaxy mass $M_{\text{gal}} = 5 \times 10^{11} M_{\odot}$, and effective radius of 7 kpc. Kick velocities were (360, 590, 830, 1070) km s^{-1} (blue, red, green, black). Open circles indicate the approximate time at which the SMBH would come to rest after “Phase I” (see text). (b) Fraction of kicked SMBHs, in a galaxy with SMBH mass M_{BH} , that remain above the torus, of radius r_{\min} , for a time of 3×10^7 yr or longer. Thick (green) contours are based on the full N -body trajectories; thin (white) contours include only “Phase I.” Gray-scale density is proportional to the fraction of kicks that result in escape, from ~ 0.3 on the left to ~ 0.05 on the right. Kicks were generated assuming “log-uniform” distributions of SMBH masses and spins (see text) with $0.3 \leq m_2/m_1 \leq 1$ and $0.5 \leq a_{1,2} \leq 0.9$. The red line shows the approximate outer radius expected for the obscuring torus (based on eqn (3), and converting M_{BH} to L assuming accretion at $0.1L_{\text{edd}}$). (c) Velocity distributions that would be observed in a representative sample of quasars, assuming that kicks occur randomly in time; additional features of the model are described in the text. The black line uses the baseline model of SMBH masses and spins in the pre-recoil binary (the same model used in Fig. 1b). Red line: like the base model, but $a = 0.3$ for both SMBHs. Blue line: equal-mass SMBHs with maximal spins. The dotted lines exclude SMBHs that are more than 10 kpc from the galaxy center at the moment of observation. The gray dashed line shows the input kick distribution for the baseline model, unmodified by motion through the galaxy.